



fraserhealth

Better health.
Best in health care.

-
- Hospice Palliative Care Program
Symptom Guidelines



Cough



Cough

□ Rationale

This guideline is adapted for inter-professional primary care providers working in various settings in Fraser Health, British Columbia and the Fraser Valley Cancer Center and any other clinical practice setting in which a user may see the guidelines as applicable.

Of patients living with and dying from advanced life threatening illness up to 86% experience the symptom of cough.^(1, 2)

□ Scope

This guideline provides recommendations for the assessment and symptom management of adult patients (age 19 years and older) living with advanced life threatening illness and experiencing the symptom of cough. This guideline does not address disease specific approaches in the management of cough. When cough becomes chronic it can be the source of sleep disturbance, agitation or anxiety and becomes exhausting. Cough is both an objective and subjective experience and has been reported that cough is multi-factorial in origin in 42% to 62% of cases. It is experienced by 23 to 86% of cancer patients and up to 59% of non-malignant patients.⁽¹⁻¹⁰⁾

□ Definition of Terms

Cough has two functions: to prevent foreign material entering the lower respiratory tract and to clear secretions from the lungs and bronchial tree.⁽³⁾

□ Standard of Care

1. Assessment
2. Diagnosis
3. Education
4. Treatment: Nonpharmacological
5. Treatment: Pharmacological

Recommendation 1 Assessment of Cough

Ongoing comprehensive assessment is the foundation of effective cough management, including interview, physical assessment, medication review, medical and surgical review, psychosocial and physical environment review (including past and present occupation) and appropriate diagnostics (*see Table 1*). Cough assessment must determine the cause, effectiveness and impact on quality of life.^(1-3, 6-8, 10-12)

Table 1: Cough Assessment using Acronym O, P, Q, R, S, T, U and V^(8, 12)

O Onset	When did it begin? How long does it last? How often does it occur?
P Provoking / Palliating	What brings it on? What makes it better? What makes it worse?
Q Quality	What does it feel like? Can you describe it? Is it positional?
R Region / Radiation	What areas are involved in your cough? Throat? Chest?
S Severity	What is the intensity of this symptom (On a scale of 0 to 10, with 0 being none and 10 being worst possible)? Right Now? At Best? At Worst? On Average? Are there any other symptom(s) that accompany this symptom?
T Treatment	What medications and treatments are you currently using? How effective are these? Do you have any side effects from the medications and treatments? What medications and treatments have you used in the past?
U Understanding / Impact on You	What do you believe is causing this symptom? How is this symptom affecting you and/or your family?
V Values	What is your goal for this symptom? What is your comfort goal or acceptable level for this symptom (On a scale of 0 to 10 with 0 being none and 10 being worst possible)? Are there any other views or feelings about this symptom that are important to you or your family?

* Physical Assessment (as appropriate for symptom)

Recommendation 2 Diagnosis

Management should include treating reversible causes where possible and desirable according to the goals of care. The most significant intervention in the management of cough is identifying underlying cause(s) and treating as appropriate. While underlying cause(s) may be evident, treatment may not be indicated, depending on the stage of the disease.

Whether or not the underlying cause(s) can be relieved or treated, all patients will benefit from management of the symptom using education or medications.

Identifying the underlying etiology of the cough is essential in determining the treatment required (*see Table 2*).^(1-3, 7-9, 11-13)

Table 2: Underlying Causes of Cough & Treatment of Choice

Underlying Cause	Treatment of Choice
Amyotrophic Lateral Sclerosis (ALS)	Scopolamine, atropine or glycopyrrolate to reduce secretions to normal and comfortable moisture levels
Bronchospasm/Bronchiectasis	Bronchodilators, antibiotics
Chronic Obstructive Pulmonary Disease (COPD)/Asthma	Conventional inhalers, nebulized drugs or saline, steroids to suppress inflammation
Congestive Heart Failure	Conventional medications to decrease excess fluid
End stage weakness	Suppress and settle with suppressant, anxiolytic, scopolamine or atropine
Gastroesophageal reflux	H2 inhibitors, proton pump inhibitor, motility agents, elevate head of bed, drain contributing ascites
Infection - Pneumonia	Prevention of aspiration. Oral antibiotics may help decrease productive cough that is disturbing sleep, or causing pain or hemoptysis
Malignant pleural effusion	Thoracentesis or pleurodesis; lying on the same side can decrease related cough
Medications	Stop offending ACE inhibitor
Post radiation lung damage	Steroids
Superior Vena Cava (SVC) obstruction	Radiotherapy, steroids
Tumour related airway irritation	Radiotherapy/brachytherapy, laser treatment, self expandable stents or steroids
Upper airway cough syndrome (Post-nasal drip) – allergies, infection, sinusitis	Nasal steroids or ipratropium. Oral antibiotics for sinusitis, expectorants (guaifenesin) or anti-histamine

Recommendation 3 Education

Cough can be distressing to experience and often more difficult to witness. Providing education regarding the etiology of cough and expectations of treatment is foundational to enhance the patient and family's ability to cope.⁽¹⁾

- As a person weakens, the ability to raise sputum is reduced.⁽¹⁾ Teach the patient 'huffing or forced expiratory technique' to clear secretions with minimal effort.^(8, 12, 14, 15)

Recommendation 4 Treatment: Nonpharmacological

- Positioning to promote and facilitate secretion drainage (postural drainage),^(1-4, 6, 8, 10, 12) but should not be used during acute exacerbation of chronic bronchitis.⁽⁸⁾
- Avoidance of smoking, chemical irritants or noxious fumes.^(4, 8-10, 12)
- Nebulized saline, steam or cold air humidifier.^(1, 3, 10-12)
- Adequate hydration.⁽²⁾
- Suction is not indicated except in the following situations: When acute fulminant pulmonary edema is present, or to clear bronchial secretions in patients with tracheostomy, or to clear saliva when full esophageal obstruction is present **or** when bleeding is present in mouth or throat.^(1, 10)
- Provide instruction in anxiety reduction strategies.⁽⁸⁾

Recommendation 5 Treatment: Pharmacological

- For minor or simple cough expectorants are effective.⁽¹⁾ Guaifenesin is a useful expectorant.
- For mild to moderate cough
 - Dextromethorphan (non-opioid) 15 to 30 mg PO q.i.d.^(2, 3, 5, 10, 12)
- For advanced disease multi-factorial cough
 - Antitussive opioids include:
 - Methadone 2.5 to 10 mg PO.⁽¹⁶⁾ Methadone is a powerful antitussive and has activity superior to those of morphine and codeine.⁽¹⁶⁾ Taking it 2 hours before bed is recommended for troublesome night cough.⁽¹⁶⁾
 - Hydromorphone 1 to 2 mg PO Q4H.^(2, 3, 5-8, 10, 17)
 - Hydrocodone 5 to 10 mg PO every 4 to 6 hours with a daily dose no higher than 60 mg.⁽¹⁶⁾ It has greater antitussive activity than codeine but less than morphine.⁽¹⁶⁾

Recommendation 5 Treatment: Pharmacological continued...

- Morphine up to 5 mg PO every 4 hours.⁽¹⁶⁾
- Codeine 10 to 20 mg PO every 4 to 6 hours, with a daily dose no higher than 120 mg.⁽¹⁶⁾
- Oxycodone 5 mg every 4 hours or 10 mg PO sustained-release oxycodone every 12 hours.⁽¹⁶⁾
- Dexamethasone 4 to 8 mg PO or I.V. or S.C. daily depending on severity and cause; taper and avoid long term use if possible (increased risk of proximal myopathy which can be very debilitating).⁽⁴⁾
- For intractable cough nebulized local anesthetics can be useful: ^(1-5, 7, 12)
 - May precipitate bronchospasm in asthmatic patients.⁽¹⁻³⁾
 - Gag reflex is inhibited after administration, keep NPO for 1 to 2 hours afterward,⁽¹⁻³⁾ and rinse and spit after nebulization to minimize numbness of lips and tongue. Use a mouthpiece rather than a mask for inhalation.
 - Bupivacaine 0.25% 5 mL q4h.⁽¹⁻³⁾
 - Lidocaine 2% 2 to 5 mL in 1 mL of normal saline q4h.⁽¹⁻³⁾
- Sodium cromoglycate has been found to be a useful cough suppressant in advanced lung cancer.^(8, 18)

□ References

Information was compiled using the CINAHL, Medline (1996 to April 2006) and Cochrane DSR, ACP Journal Club, DARE and CCTR databases, limiting to reviews/systematic reviews, clinical trials, case studies and guidelines/protocols using respiratory/cough terms in conjunction with palliative/hospice/end of life/dying. Palliative care textbooks mentioned in generated articles were hand searched. Articles not written in English were excluded.

1. Gallagher R. Cough in Terminal Care. In: Downing GM, Wainwright W, editors. *Medical Care of the Dying*. 4th ed. Victoria, B.C. Canada: Victoria Hospice Society Learning Centre for Palliative Care; 2006. p. 363 - 5.
2. Chan KS, Sham M, M. K., Tse DMW, Thorsen AB. Palliative medicine in malignant respiratory diseases. In: Doyle D, Hanks G, Cherny NI, Calman K, editors. *Oxford Textbook of Palliative Medicine*. 3rd ed. New York, New York: Oxford University Press Inc., New York; 2005. p. 587 - 618.
3. Leach R. Palliative medicine and non-malignant, end-stage respiratory disease. In: Doyle D, Hanks G, Cherny NI, Calman K, editors. *Oxford Textbook of Palliative Medicine*. 3rd ed. New York, New York: Oxford University Press Inc., New York; 2005. p. 895 - 916.
4. Esper P, Heidrich D. Symptom Clusters in Advanced Illness. *Seminars in Oncology Nursing*. 2005 February 2005;21(1):20 - 8.
5. Kvale PA, Simoff M, Prakash UBS. Palliative care. *Chest* 2003 Jan; 123(1): Suppl: 284S-311S. January 2003.
6. Louie K, Bertolino M, Fainsinger R. Management of Intractable Cough. *Journal of Palliative Care*. 1992 1992;8(4):46 - 8.
7. Zyllicz Z, Krajnik M. The use of antitussive drugs in terminally ill patients. *European Journal of Palliative Care* 2004 Nov-Dec; 11(6): 225-9.
8. Irwin RS, Baumann MH, Bolser DC, Boulet L-P, Braman SS, Brightling CE, et al. Diagnosis and Management of Cough - Executive Summary. *Chest*. 2006 January 2006;129:1S - 23S.
9. Holmes RL, Fadden CT. Evaluation of the Patient with Chronic Cough. *American Family Physician*. 2004 May 1, 2004;69(9):2159 - 66.
10. Whiting N. Cough. In: MacDonald N, Oneschuk D, Hagen N, Doyle D, editors. *Palliative Medicine - A case based manual* 2nd ed. New York: Oxford University Press Inc.; 2005.
11. Sherman DW, Matzo ML, Coyne P, Ferrell BR, Penn BK. Teaching symptom management in end-of-life care: the didactic content and teaching strategies based on the end-of-life nursing education curriculum. *Journal for Nurses in Staff Development - JNSD*. 2004 May-Jun;20(3):103-15; quiz 16-7.
12. Coyne PJ, Lyne ME, Watson AC. Symptom management in people with AIDS. *American Journal of Nursing*. 2002 Sep;102(9):48-56; quiz 7.
13. Dudgeon D. Dyspnoea. In: MacDonald N, Oneschuk D, Hagen N, Doyle D, editors. *Palliative Medicine - A case based manual* 2nd ed. New York: Oxford University Press Inc.; 2005.
14. McCool FD, Rosen MJ. Nonpharmacologic Airway Clearance Therapies: ACCP Evidence-Based Clinical Practice Guidelines. *Chest*. 2006 January, 2006;129(Supplement):250 - 9.
15. Registered Nurses' Association of Ontario. Nursing Best Practice Guideline: Nursing Care of Dyspnea: The 6th Vital Sign in Individuals with Chronic Obstructive Pulmonary Disease (COPD). Nursing Best Practice Guidelines Program 2005; Available from: http://www.rnao.org/Storage/11/604_BPG_COPD.pdf
16. Homsy J, Walsh D, Nelson KA. Important drugs for cough in advanced cancer. *Support Care Cancer*. 2001;19:565-74.
17. Stein WM, Young KM. Nebulized morphine for paroxysmal cough and dyspnea in a nursing home resident with metastatic cancer. *The American Journal of Hospice and Palliative Care*. 1997 March/April 1997;14(2):52 - 6.
18. Moroni M, Porta C, Gualtieri G, Nastasi G, Tinelli C. Inhaled sodium cromoglycate to treat advanced lung cancer patients. *British Journal of Cancer*. 1996;74(2):309-11.

Approved by: Hospice Palliative Care, Clinical Practice Committee, November 24, 2006