### **During the Appointment**

Video or Electronystagmography (VNG/ENG) is an assessment used to help determine whether your balance system is functioning within the range of normal.

A comprehensive battery of tests will be performed during your appointment. Prior to each test, an explanation will be given so you will have a better understanding of what is being tested and why. We make every attempt to make your visit comfortable as well as educational.

Portions of this test might cause a brief sensation of motion, but this tends to be mild and lasts from a few seconds to a couple of minutes.

If possible, we encourage you to have someone accompany you to and from the appointment. However, if this is not possible, try to plan your day to include an extra 15 to 30 minutes to rest after your test and before leaving the hospital.

Your VNG/ENG ap	pointment
is scheduled	on:

Name: \_\_\_\_\_

Date: \_\_\_\_\_

Location:

#### Surrey Memorial Hospital Rehabilitation Services 13750 - 96<sup>th</sup> Ave Surrey BC

Abbotsford Regional Hospital Rehabilitation Services 32900 Marshall Rd Abbotsford BC

#### Please Note:

- We require a <u>minimum of 48</u> <u>hours</u> notice to cancel or reschedule your appointment.
- If you are more than 15 minutes late for your appointment, it may not be possible to complete the test. You would need to return another day to do so.



## VNG / ENG (Video/Electronystagmography)

Pre-Test Instructions And Assessment Information

Acute Audiology Services Fraser Health Authority

Phone: 604-585-5674 Fax: 604-585-5568

# Pre-Test Instructions

Certain medications can influence the body's response to the test, thus giving a false or misleading result. Therefore, for <u>at least 48 hours prior to the</u> <u>test</u>, please do not take the following:

- Alcohol in any form, including cough medications containing alcohol
- Caffeine coffee, tea, pop, chocolate
- Herbal remedies e.g., Gingko Biloba, Valerian
- Antihistamines found in many cold and allergy medications. e.g., Chlortriplon, Claritin, Benedryl, Triaminic
- Anti-nausea medication e.g. Gravol, Dramamine, Penergan, Compazine, Bonine, Transdermal
- Anti-vertigo medications E.g., Antivert, Serc (novo-betahistine), Benedryl, Meclizine
- Stimulants E.g., amphetamines, appetite suppressants, "stay awake" medications such as No Doz
- Anti-seizure medications (when not taken for Epilepsy) – E.g., Dilantin, Tegretol
- Diuretics / "water pill" E.g., Dyazinde, Lasix, Neptazane

## Pre-Test Instructions (Continued)

- Pain medications including migraine meds, analgesics, muscle relaxants, narcotics, and barbiturates (e.g., Codeine, Demerol, Phenaphen, Phenobarbital, Tylenol, Ibuprofen, Aspirin products, Amitriptyline, Proparanolol)
- Sedatives including anti-anxiety medications, sleeping pills, and tranquilizers (e.g., Tylenol PM, Seconal, Restoril, Serax, Librium, Halcion, Ativan, Lorazepam, Zoloft, Prozac, Xanax, Diazepam)
- Tricyclic anti-depressants -- E.g., Elavil, Amitriptyline, Amoxapine, Wellbutrin
- Recreational drugs of any kind
- Methadone
- Quinine

If you have concerns whether or not to take a particular medication, please consult with your referring physician to determine if you can safely stop the medication for the 48 hours or if you can use an alternative medication that is not restricted.

You should always consult with your physician before discontinuing any prescribed medication.

# The following medications **should** be taken as usual:

Antibiotics Blood pressure pills (aside from diuretics) Heart medications Thyroid pills Diabetes pills and insulin Epilepsy control Antipsychotics Hormone replacement Birth control

For at least <u>2 hours</u> prior to the test, please also:

#### Do not eat anything Do not drink anything Do not smoke

- Bring a list of all your medications
- Please inform us if you have any neck/back pain.
- Do not wear make-up or mascara.
- Dress comfortably.
- Bring your questionnaire with you.