


Donning PPE: Droplet Precautions and Aerosol Generating Procedures for COVID-19

- Assessment, treatment and testing of suspected or confirmed COVID-19 patients require Droplets Precautions Personal Protective Equipment (PPE).
- This PPE is required for all entry into rooms with suspected or confirmed COVID-19 patient.
- It includes a gown, surgical/procedure mask, protective eyewear or face shield and gloves.
- Use an N95 respirator if performing an Aerosol Generating Procedure (AGP).



1. Gather required materials




Rub hands until dry ensuring all surfaces of the hand are covered.

2. Perform hand hygiene



Tie neck and waist ties securely.

3. Put on gown




Place mask over nose and under chin, secure ties, loops and straps. Mould metal piece to the bridge of your nose. Perform seal check if applicable.

4. Put on a surgical/procedure mask (or an N95 respirator for AGPs)



5. Put on protective eyewear or a face shield (for AGPs, wear eye goggles or face shield)



Pull the gloves completely over the cuff of the gown sleeve.

6. Put on gloves

Doffing PPE: Droplet Precautions and Aerosol Generating Procedures for COVID-19

- Assessment, treatment and testing of suspected or confirmed COVID-19 patients require Droplet Precautions Personal Protective Equipment (PPE).
- This PPE is required for all entry into rooms with suspected or confirmed COVID-19 patient.
- It includes a gown, surgical/procedure mask, protective eyewear or face shield and gloves.
- Use an N95 respirator if performing an Aerosol Generating Procedure (AGP).


PPE must be removed at least 2 meters away from the patient




1. Remove gloves




2. Perform hand hygiene



3. Remove gown



4. Perform hand hygiene



5. Remove eye protection or face shield (if re-using face shield, remove outside the patient room)



6. Perform hand hygiene



7. Remove surgical/procedure mask or N95 respirator



8. Perform hand hygiene